

Use of targeted muscle reinnervation to reduce phantom limb pain in people with amputation

Jennifer Hall - MPO Student - jhall12@uw.edu

University of Washington - School of Rehabilitation Medicine - Division of Prosthetics and Orthotics

Creation date: March 2019

References

1. Valerio IL, Dumanian GA, Jordan SW, et al. Preemptive Treatment of Phantom and Residual Limb Pain with Targeted Muscle Reinnervation at the Time of Major Limb Amputation. *J Am Coll Surg*. 2019;228(3):217-226. doi:10.1016/j.jamcollsurg.2018.12.015
2. Dumanian GA, Potter BK, Mioton LM, et al. Targeted Muscle Reinnervation Treats Neuroma and Phantom Pain in Major Limb Amputees: A Randomized Clinical Trial. *Ann Surg*. doi:10.1097/SLA.0000000000003088
3. Kuiken TA, Barlow AK, Hargrove L, Dumanian GA. Targeted Muscle Reinnervation for the Upper and Lower Extremity. *Tech Orthop Rockv Md*. 2017;32(2):109-116. doi:10.1097/BTO.000000000000194
4. Bowen JB, Wee CE, Kalik J, Valerio IL. Targeted Muscle Reinnervation to Improve Pain, Prosthetic Tolerance, and Bioprosthetic Outcomes in the Amputee. *Adv Wound Care*. 2017;6(8):261-267. doi:10.1089/wound.2016.0717
5. Bowen JB, Ruter D, Wee C, West J, Valerio IL. Targeted Muscle Reinnervation Technique in Below-Knee Amputation: *Plast Reconstr Surg*. 2019;143(1):309-312. doi:10.1097/PRS.0000000000005133