

## References

1. Suleman F, Sohail A, Javed G, Samar SS. The role of helmet therapy in craniosynostosis: a systematic review. *Asian J Neurosurg*. 2024;19(3):610-617. <https://doi.org/10.1055/s-0044-1791228>
2. Riordan CP, Zurakowski D, Meier PM, et al. Minimally invasive endoscopic surgery for infantile craniosynostosis: a longitudinal cohort study. *J Pediatr*. 2020;216:142-149.e2. <https://doi.org/10.1016/j.jpeds.2019.09.037>
3. Nguyen HT, Washington GN, Cepeda A, et al. Evaluation of helmeting therapy duration after endoscopic strip craniectomy for metopic and sagittal craniosynostosis. *J Craniofac Surg*. 2024;35(2):415-418. <https://doi.org/10.1097/SCS.00000000000009887>
4. Cross C, Delye H, Khonsari RH, Moazen M. A preliminary analysis of replicating the biomechanics of helmet therapy for sagittal craniosynostosis. *Childs Nerv Syst*. 2023;39(4):989-996. <https://doi.org/10.1007/s00381-022-05792-1>
5. Iyer RR, Ye X, Jin Q, et al. Optimal duration of postoperative helmet therapy following endoscopic strip craniectomy for sagittal craniosynostosis. *J Neurosurg Pediatr*. 2018;22(6):610-615. <https://doi.org/10.3171/2018.5.PEDS184>
6. Rudolph CM, Awad AN, Toth K, Adamo MA, Carpenter C. Endoscopic strip craniectomy and helmet therapy for sagittal craniosynostosis: an analysis of cranial growth changes in the early postoperative period. *Cleft Palate Craniofac J*. 2024;61(3):458-468. <https://doi.org/10.1177/10556656221131119>