

A polycentric hydraulic hip joint increases prosthetic gait biomechanics and functional capabilities for hip disarticulation amputees compared to the current standard of care

R. Jenkins, CPO; Bionic Prosthetics and Orthotics; rljenkins716@gmail.com

Creation Date: April 2023, Date of Reassessment: April 2028

References:

1. Nelson LF, Carbone NT. Functional outcome measurements of a veteran with a hip disarticulation using a helix 3D hip joint: a case report. *Journal of Prosthetics and Orthotics (JPO)*. 2011; 23(1), 21-26.
2. Ludwigs E, Kannenberg A, Wüstefeld D. Evaluation of the benefits of a new prosthetic hip joint system in activities of daily function in patients after hip disarticulation or hemipelvectomy. *Journal of Prosthetics and Orthotics (JPO)*. 2013; 25(3), 118-126.
3. Ludwigs E, Bellmann M, Schmalz T, Blumentritt S. (2010). Biomechanical differences between two exoprosthetic hip joint systems during level walking. *Prosthetics and Orthotics International*. 2010; 34(4), 449–460.
4. Karimi M, Kamali M, Omar H, Mostman J. Evaluation of gait performance in a hemipelvectomy amputation walking with a Canadian prosthesis. *Case Reports in Orthopedics*. 2014; 2014 (Article ID 962980), 1-8.
5. Gailledrat E, Moineau B, Seetha V, DeAngelis MP, Saurel B, Chabloz P, Nougier V, Pérennou D. Does the new Helix 3D hip joint improve walking of hip disarticulated amputees?. *Annals of Physical and Rehabilitation Medicine*. 2013; 56(5), 411–418.
6. Kawaguchi T, Yamada T, Iwashita K. Biomechanical gait analysis for a hip disarticulation prosthesis: power source for the swing phase of a hip disarticulation prosthetic limb. *Journal of Physical Therapy Science*. 2023; 35(5), 361-365.

