

Cardiovascular System - Common Medications

Diuretics: "water pills"

Uses: hypertension, edema, heart failure

Side effects: Side effects: symptoms of low blood pressure, frequent urination, change in electrolytes

Subclass	Generic (Brand)	Available Formulations
Thiazides and thiazide-like	Hydrochlorothiazide(Microzide)	Tablet, capsule
	Chlorthalidone (Diuril)	Tablet
	Chlorothiazide (Diuril)	Tablet, suspension
	Metolazone (Zaroxolyn)	Tablet
Loops	Furosemide (Lasix)	Tablet, solution, injection
	Bumetanide (Bumex)	Tablet, injection
	Torsemide (Demaxex)	Tablet, IV
Potassium Sparing	Spironolactone (Aldactone)	Tablet
	Eplerenone (Inspra)	Tablet
	Triamterene (Dyrenium)	Capsule

Angiotensin Converting Enzyme (ACE) Inhibitors: "-pril"

Uses: hypertension, heart failure, and to increase survival after heart attack

Side Effects: dry cough, increase potassium, reduced kidney function, angioedema, anaphylaxis, symptoms of low blood pressure

DDIs: Potassium supplements, NSAIDs

Pregnancy Category: X

Generic (Brand)	Available Formulations
Benazepril (Lotensin)	Tablet
Captopril (Capoten)	Tablet
Enalapril (Vasotect)	Tablet, powder for solution
Lisinopril (Prinivil, Zestril)	Tablet
Quinapril (Accupril)	Tablet
Ramipril (Altace)	Capsule

Angiotension Receptor Blockers (ARBs): "-sartan"

Uses: hypertension, heart failure

Side Effects: Increase potassium, symptoms of low blood pressure, reduced kidney function, anaphylaxis

DDIs: K+ supplements

Pregnancy Category: X

Generic (Brand)	Available Formulations
Losartan (Cozaar)	Tablet
Valsartan (Diovan)	Tablet
Olmesartan (Benicar)	Tablet
Candesartan (Atacand)	Tablet
Irbesartan (Avapro)	Tablet
Telmisartan (Micardis)	Tablet

Calcium Channel Blockers (CCBs)

Uses: Hypertension, non-dihydropyridines can also be used for dysrhythmias

Side Effects: flushing, headache, swelling of feet and ankles, symptoms of low blood pressure.

Drug-food interactions: avoid grapefruit juice while taking non-dihydropyridines

Subclass	Generic (Brand)	Available Formulations
Dihydropyridine: "pine"	Amlodipine (Norvasc)	Tablet
	Nifedipine (Adalat CC, Procardia XL)	Capsule, ER tablet
	Nicardipine (Cardene, Cardene SR)	Capsule, ER capsule, IV
	Felodipine (Plendil ER)	ER tablet
Non-dihydropyridine	Diltiazem (Cardizem, Dilacor XR, etc.)	Tablet, ER Tablet, ER capsule, IV, powder for IV solution
	Verapamil (Calan, Calan SR)	Tablet, ER tablet, ER capsule, IV

Beta Blockers: “-lol”

Uses: Heart failure, hypertension, increase survival after heart attack, some used for dysrhythmias
Side Effects: slow heart rate, fatigue, decreased ability to exercise, symptoms of low blood pressure
Drug-disease interactions: lung conditions (asthma, emphysema, etc.), diabetes, depression

Generic (Brand)	Available Formulations
Atenolol (Tenormin)	Tablet
Bisoprolol (Zebeta)	Tablet
Metoprolol (Lopressor, Toprol XL)	Tablet, IV, ER tablet
Nebivolol (Bystolic)	Tablet
Propanolol (Inderal LA, Inderal XL)	Tablet, ER capsule, solution, IV
Carvedilol (Coreg)	Tablet, ER tablet
Labetalol (Trandate)	Tablet, injection, IV

Vasodilators

Uses: when other therapies don't work and in emergency situations

Side Effects: fast heart rate, swelling

Generic (Brand)	Available Formulations
Hydralazine (Apresoline)	Tablet, injection
Sodium Nitroprusside (Nitropress)	IV, powder for IV solution
Minoxidil (Loniten)	Tablet, topical solution

Individual Agents

Generic (Brand)	Available Formulations
Clonidine (Catapres): Class: alpha 2 agonist <u>Uses:</u> hypertension after failure with other classes <u>Side Effects:</u> dry mouth, drowsiness, fatigue, skin irritation from patch, rebound hypertension if discontinued abruptly	Tablet, ER tablet, patch
Digoxin (Lanoxin): <u>Uses:</u> relieve symptoms and decrease hospitalizations in heart failure patients, anti-dysrhythmic agent <u>Drug-disease interactions:</u> avoid in renal disease <u>Other precautions:</u> use with caution in elderly	Tablet, solution, injection, or IV

HMG-CoA reductase inhibitors: “-statins”

Uses: First line for hyperlipidemia, prevention of heart attacks

Side Effects: generally well tolerated, can cause mild to severe mm pain, rhabdomyolysis (severe but rare), irritation to the liver

Drug food interactions: avoid grapefruit juice

Generic (Brand)	Available Formulations
Atorvastatin (Lipitor)	Tablet
Simvastatin (Zocor)	Tablet
Lovastatin (Mevacor)	Tablet
Pravastatin (Pravachol)	Tablet
Rosuvastatin (Crestor)	Tablet
Fluvastatin (Lescol, Lescol XL)	Capsule, ER tablet

Fibrates

Uses: decrease triglycerides and increase HDL levels

Side Effects: headache, nausea, vomiting, diarrhea, abdominal pain, rash, liver irritation, increased LDL

Generic (Brand)	Available Formulations
Fenofibrate (Tricor)	Tablet, capsule
Gemfibrozil (Lopid)	Tablet

Bile Acid Sequestrants

Uses: decrease LDL, not commonly used due to intolerable side effects for many patients

Side Effects: nausea, bloating, constipation, flatulence, increase triglycerides

DDIs: bind to many drugs and disrupt absorption, so take at a different time of day

Generic (Brand)	Available Formulations
Cholestyramine (Questran)	Powder for suspension
Colesevelam (WelChol)	Tablet, powder for suspension
Colestipol (Colestid)	Tablet, powder for suspension

Niacin (Vitamin B3)	
Generic (Brand)	Available Formulations
Niacin (Niacor, Niaspan) <u>Uses:</u> Second line choice for hyperlipidemia (decreases LDL and triglycerides, increases HDL) <u>Side Effects:</u> intense flushing, itching, liver irritation <u>Management of intense flushing:</u> Take acetaminophen or an NSAID 30 minutes prior. If necessary, can switch to ER formulation, which is not associated with flushing.	Niacin IR capsule and IR tablet, OTC Niacor Niacin ER capsule and ER tablet, Slo-Niacin OTC, Niaspan

Anti-dysrhythmic drugs	Generic (Brand)	Available Formulations
<u>Uses:</u> arrhythmia <u>Side Effects:</u> can cause new or more life threatening dysrhythmias	Amiodarone (Pacerone)	Tablet, IV
	Dronedarone (Multaq)	Tablet
	<u>Pregnancy Category: X</u>	
	Dofetilide (Tikosyn) REMS, must be started as an inpatient	Capsule
	Sotalol (Betapace)	Tablet, IV
	Diltiazem	*See CCBs
	Verapamil	*See CCBs