


High level areas to address when considering resumption of business. More detailed information for each topic can be found on pages 2-4.



- Request that all employees wear a mask when possible.
- If you do not have surgical masks to provide, encourage employees to bring cloth masks from home.

5/5/20

Masks



Communication

	Policies It is important that your employees know what you are doing to keep them safe and what will be required of them. Consider developing and distributing clear policies for the following:		
•			
		Remote work	
		☐ Symptom screening	
			Door screens
			Self-monitoring
		☐ Testing	
		☐ Return to work	
			When to stay home or self-isolate
			When to come back to work
		☐ Notification and tracking of symptoms or test results	
	☐ Pay and Leave		
		□ Social distancing – employees and patrons□ Cleaning and Sanitizing□ Masks	
•	Basic guidelines for most of the above are listed on pages 3-4, but they are not comprehensive and should be used as a base for more detailed policies or standard operating procedures. OSHA's COVID-19 page should be consulted for more information about how to protect your employees. CDC's Interim Guidance for Business and Employers to Plan and Respond to Coronavirus Disease 2019 (COVID-19) should be consulted for additional resources. Your state may have additional requirements to adhere to when resuming business.		
	Education		
•	Consider providing your employees with resources so that they know how to keep themselves		

- - Resources could include:
 Video trainings
 - Email series around workplace safety during COVID-19
 - Virtual town hall meetings
 - An employee resource phone line

and others safe, both at work and at home.

5/5/20 2



3

Screening and Testing

■ Symptom and Temperature Screening

- Develop processes for implementing screenings for <u>symptoms of COVID-19</u>, including fever
- · We suggest this be done via self monitoring and screening at the door before staff enter the facility
- Employees experiencing symptoms of COVID-19 should self-isolate according to <u>CDC guidelines</u> and contact their healthcare provider.

<u>Self Monitoring</u>: encourage employees to take their temperature twice daily, including once immediately before starting their shifts or coming to work, and track other symptoms. If everything is normal, they may come to work. If they have a fever of 100 degrees Fahrenheit or higher, cough, shortness of breath, or any of the other symptoms of COVID-19, they should stay home and notify their manager.

<u>Door Screenings</u>: if you have the staff and Personal Protective Equipment, you may be able to operationalize screenings for temperature <u>and</u> symptoms at the door. This will require thermometers, thermometer covers, masks, gloves, and training of your staff to be screeners.

□ Testing

- COVID-19 diagnostic testing is primarily indicated for individuals who are symptomatic (cough, shortness of breath or difficulty breathing, or two or more of the following: fever, chills, repeated shaking with chills, muscle pain, headache, sore throat, new loss of taste or smell).
- Symptomatic employees should self-isolate and be referred to a healthcare provider for testing.
 - Piedmont has several ways to access care for COVID related evaluations, including telemedicine and virtual urgent care visits. Our healthcare providers can help employees navigate testing, treatment and return-to-work determinations quickly and efficiently.

☐ Return to Work Guidance

- <u>The CDC has clear guidelines</u> outlining when an individual should discontinue self isolation following COVID-19 symptoms or a positive COVID-19 test. *Please note that these may change and that there are separate guidelines for healthcare workers:*
- Symptom-based strategy: Individuals self-isolating due to COVID-19 symptoms may stop isolating once:
 - At least 3 days (72 hours) have passed since recovery defined as resolution of fever without
 the use of fever-reducing medications and improvement in respiratory symptoms (e.g.,
 cough, shortness of breath); and,
 - At least 10 days have passed since symptoms first appeared.
- Test-based strategy: may be difficult to operationalize without readily available test supply.
 - Resolution of fever without the use of fever-reducing medications and
 - Improvement in respiratory symptoms (e.g., cough, shortness of breath), and
 - At least two consecutive negative test results collected ≥24 hours apart

We encourage you to consider the potential impact that the self-isolation requirements may have on staffing and pay/leave policies. Developing a clear process to track employees that are self-isolating due to symptoms or positive test results may be helpful.



Workplace Safety and Infection Prevention

■ Social Distancing

- Consider flexible work hours or schedules; utilizing remote options whenever possible
- Space chairs and desks at least 6 feet apart in all directions
- Use tape lines on floor to indicate 6 ft. marks
- Declutter desks, break rooms and conference rooms to remove non-essential items from countertops
- Discuss options for limiting conference room use or spreading out inside conference rooms
- Waiting rooms should be devoid of "hi-touch" items such as magazines, reading material, candy bowls and coffee machines
- Use plexiglass between staff and patrons, if available and where possible
- Work with building management to develop guidelines for elevators and coordinate other requirements for social distancing

□ Cleaning

- Ensure that you or your cleaning service have an appropriate supply of cleaning and disinfecting products that are <u>validated for use against COVID-19</u>
- Work with your cleaning service to develop a schedule for more frequent cleaning of restrooms and high-touch surfaces
- Encourage employees to practice hand hygiene and keep their desks and personal belongings clean
- Discourage employees from using coworkers' phones, desks, or work tools when possible

□ Masks

- CDC recommends wearing cloth face coverings in public settings where other social distancing measures are difficult to maintain to slow the spread of the virus
- Encourage employees to wear masks as frequently as possible, especially when using common spaces in the workplace like restrooms, conference rooms and breakrooms
 - If you do not have surgical masks to provide, <u>homemade cloth face coverings</u> are sufficient in most settings
 - Be sure to check the recommended PPE for your <u>job risk classification</u> when developing guidelines around masking

5/5/20